

**881.** Proposed by Kim McInturff, Santa Barbara, California.

Prove that for every positive integer  $n$

$$\sum_{k=0}^{n-1} \binom{2k}{k} = \sum_{i=1}^n \frac{2 \sin \frac{2i\pi}{3}}{\sqrt{3}} \binom{2n}{n+i}.$$

**Solution:** (by José M. Pacheco and Ángel Plaza, University of Las Palmas de Gran Canaria, 35017-Las Palmas G.C., Spain)

First, notice that the coefficient  $\frac{2}{\sqrt{3}} \sin \left( \frac{2i\pi}{3} \right)$  takes the values  $+1, -1, 0, +1, -1, 0, \dots$ , and thus the right-hand side (RHS) may be written as  $\sum_{i=1}^n i_{[3]} \binom{2n}{n+i}$ , where  $i_{[3]}$  represents, for  $i = 1, 2, \dots$ , the successive values of the sequence  $+1, -1, 0, +1, -1, 0, \dots$

The equation may be proved by induction. For the first values of  $n$  we have:

- $n = 1$ :  $\binom{0}{0} = +\binom{2}{2} = 1$
- $n = 2$ :  $\binom{0}{0} + \binom{2}{1} = +\binom{4}{2+1} - \binom{4}{2+2} = 3$
- $n = 3$ :  $\binom{0}{0} + \binom{2}{1} + \binom{4}{2} = +\binom{6}{3+1} - \binom{6}{3+2} + 0\binom{6}{3+3} = 9$

Let us suppose the expression is true for  $n$ . Now, for  $n + 1$  it is:

$$(LHS) = \sum_{k=0}^n \binom{2k}{k} = \binom{2n}{n} + \sum_{k=0}^{n-1} \binom{2k}{k} \quad (1)$$

$$(RHS) = \sum_{i=1}^{n+1} i_{[3]} \binom{2n+2}{n+1+i} \quad (2)$$

By applying twice the Pascal's rule,  $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$ , then for every integer  $m$ ,  $\binom{2n+2}{m} = \binom{2n}{m-2} + 2\binom{2n}{m-1} + \binom{2n}{m}$ , and using this into Eq. (2) we have:

$$\begin{aligned}
(RHS) &= \binom{2n}{n} + \overbrace{2\binom{2n}{n+1}} + \underbrace{\binom{2n}{n+2}} \\
&\quad - \overbrace{\binom{2n}{n+1}} - \underbrace{2\binom{2n}{n+2}} - \cancel{\binom{2n}{n+3}} + 0 + \\
&\quad + \cancel{\binom{2n}{n+3}} + 2\binom{2n}{n+4} + \binom{2n}{n+5} - \dots \\
&\quad + (n+1)_{[3]} \left[ \binom{2n}{2n} + \binom{2n}{2n+1} + \binom{2n}{2n+2} \right] = \\
&= \binom{2n}{n} + \binom{2n}{n+1} - \binom{2n}{n+2} + \binom{2n}{n+4} - \binom{2n}{n+5} + \\
&\quad \dots + (n)_{[3]} \binom{2n}{2n} + c_{n+1} \binom{2n}{2n+1} + c_{n+2} \binom{2n}{2n+2}
\end{aligned}$$

It is easy to see that the coefficient of the term  $\binom{2n}{2n}$  is always  $(n)_{[3]}$ . On the other hand, the coefficients of the last two terms, noted by  $c_{n+1}$  and  $c_{n+2}$ , depend on the value of  $(n+1)_{[3]}$ , but these two terms are zero because the lower index is greater than the upper one.

Therefore,  $(RHS)$  is the sum of  $\binom{2n}{n} + \sum_{i=1}^n i_{[3]} \binom{2n}{n+i}$ , and by the induction hypothesis  $(RHS) = \binom{2n}{n} + \sum_{k=0}^{n-1} \binom{2k}{k} = \sum_{k=0}^n \binom{2k}{k} = (LHS)$  and the proof is done.  $\square$